

COSMOS Photometric Redshifts with 30-bands for 2-deg² ¹

O. Ilbert², P. Capak^{3,4}, M. Salvato³, H. Aussel⁵, H. J. McCracken⁶, D. B. Sanders², N. Scoville³, J. Kartaltepe², S. Arnouts⁷, E. Le Floc'h², B. Mobasher⁸, Y. Taniguchi⁹, F. Lamareille¹⁰, A. Leauthaud¹¹, S. Sasaki^{12,13}, D. Thompson^{3,14}, M. Zamojski³, G. Zamorani¹⁵, S. Bardelli¹⁵, M. Bolzonella¹⁵, A. Bongiorno¹¹, M. Brusa¹⁶, K.I. Caputi¹⁷, C.M. Carollo¹⁷, T. Contini¹⁰, R. Cook³, G. Coppa¹⁵, O. Cucciati¹⁸, S. de la Torre¹², L. de Ravel¹², P. Franzetti¹⁹, B. Garilli¹⁹, G. Hasinger¹⁶, A. Iovino¹⁸, P. Kampeczyk¹⁷, J.-P. Kneib¹¹, C. Knobel¹⁷, K. Kovac¹⁷, J.F. Le Borgne¹⁰, V. Le Brun¹¹, O. Le Fèvre¹², S. Lilly¹⁷, D. Looper², C. Maier¹⁷, V. Mainieri¹⁰, Y. Mellier⁶, M. Mignoli¹⁵, T. Murayama,¹³ R. Pellò¹⁰, Y. Peng¹⁷, E. Pérez-Montero¹⁰, A. Renzini²⁰, E. Ricciardelli²⁰, D. Schiminovich²¹, M. Scodeggio¹⁹, Y. Shioya⁹, J. Silverman¹⁷, J. Surace⁴, M. Tanaka²², L. Tasca¹¹, L. Tresse¹¹, D. Vergani¹⁹, E. Zucca¹⁵

ABSTRACT

We present accurate photometric redshifts in the 2-deg² COSMOS field. The redshifts are computed with 30 broad, intermediate, and narrow bands covering the UV (*GALEX*), Visible-NIR (Subaru, CFHT, UKIRT and NOAO) and mid-IR (*Spitzer*/IRAC). A χ^2 template-fitting method (*Le Phare*) was used and calibrated with large spectroscopic samples from VLT-VIMOS and Keck-DEIMOS. We develop and implement a new method which accounts for the contributions from emission lines ([O II], H β , H α and Ly α) to the spectral energy distributions (SEDs). The treatment of emission lines improves the photo- z accuracy by a factor of 2.5. Comparison of the derived photo- z with 4148 spectroscopic redshifts (i.e. $\Delta z = z_s - z_p$) indicates a dispersion of $\sigma_{\Delta z/(1+z_s)} = 0.007$ at $i_{AB}^+ < 22.5$, a factor of 2 – 6 times more accurate than earlier photo- z in the COSMOS, CFHTLS and COMBO-17 survey fields. At fainter magnitudes $i_{AB}^+ < 24$ and $z < 1.25$, the accuracy is $\sigma_{\Delta z/(1+z_s)} = 0.012$. The deep NIR and IRAC coverage enables the photo- z to be extended to $z \sim 2$ albeit with a lower accuracy ($\sigma_{\Delta z/(1+z_s)} = 0.06$ at $i_{AB}^+ \sim 24$). The redshift distribution of large magnitude-selected samples is derived and the median redshift is found to range from $z_m = 0.66$ at $22 < i_{AB}^+ < 22.5$ to $z_m = 1.06$ at $24.5 < i_{AB}^+ < 25$. At $i_{AB}^+ < 26.0$, the multi-wavelength COSMOS catalog includes approximately 607,617 objects. The COSMOS-30 photo- z enable the full exploitation of this survey for studies of galaxy and large scale structure evolution at high redshift.

Subject headings: galaxies: distances and redshifts — galaxies: evolution — galaxies: formation

¹Based on observations with the NASA/ESA *Hubble Space Telescope*, obtained at the Space Telescope Science Institute, which is operated by AURA Inc. under NASA contract NAS 5-26555. Also based on observations made with the *Spitzer Space Telescope*, which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under NASA contract 1407. Also based on data collected at : the Subaru Telescope, which is operated by the

National Astronomical Observatory of Japan; the XMM-Newton, an ESA science mission with instruments and contributions directly funded by ESA Member States and NASA; the European Southern Observatory under Large Program 175.A-0839, Chile; Kitt Peak National Observatory, Cerro Tololo Inter-American Observatory and the National Optical Astronomy Observatory, which are operated by the Association of Universities for Research in As-

1. Introduction

Photometric redshifts (hereafter photo-z) are an estimate of galaxy distances based on the ob-

tronomy, Inc. (AURA) under cooperative agreement with the National Science Foundation; and the Canada-France-Hawaii Telescope with MegaPrime/MegaCam operated as a joint project by the CFHT Corporation, CEA/DAPNIA, the NRC and CADC of Canada, the CNRS of France, TERAPIX and the Univ. of Hawaii.

²Institute for Astronomy, 2680 Woodlawn Dr., University of Hawaii, Honolulu, Hawaii, 96822

³California Institute of Technology, MC 105-24, 1200 East California Boulevard, Pasadena, CA 91125

⁴Spitzer Science Center, California Institute of Technology, Pasadena, CA 91125

⁵AIM Unité Mixte de Recherche CEA CNRS Université Paris VII UMR n158

⁶Institut d'Astrophysique de Paris, UMR7095 CNRS, Université Pierre et Marie Curie, 98 bis Boulevard Arago, 75014 Paris, France

⁷Canada France Hawaii telescope corporation, 65-1238 Mamalahoa Hwy, Kamuela, Hawaii 96743, USA

⁸Department of Physics and Astronomy, University of California, Riverside, CA, 92521, USA

⁹Research Center for Space and Cosmic Evolution, Ehime University, Bunkyo-cho, Matsuyama 790-8577, Japan

¹⁰Max Planck Institut für Extraterrestrische Physik, D-85478 Garching, Germany

¹¹Laboratoire d'Astrophysique de Toulouse/Tarbes, Université de Toulouse, CNRS, 14 avenue E. Belin, 31400 Toulouse, France

¹²Laboratoire d'Astrophysique de Marseille, BP 8, Traverse du Siphon, 13376 Marseille Cedex 12, France

¹³Astronomical Institute, Graduate School of Science, Tohoku University, Aramaki, Aoba, Sendai 980-8578, Japan

¹⁴Physics Department, Graduate School of Science & Engineering, Ehime University, 2-5 Bunkyo-cho, Matsuyama, 790-8577, Japan

¹⁵LBT Observatory, University of Arizona, 933 N. Cherry Ave., Tucson, Arizona, 85721-0065, USA

¹⁶INAF-Osservatorio Astronomico di Bologna, via Ranzani 1, I-40127 Bologna, Italy

¹⁷Department of Physics, ETH Zurich, CH-8093 Zurich, Switzerland

¹⁸INAF Osservatorio Astronomico di Brera, Milano, Italy

¹⁹INAF - IASF Milano, via Bassini 15, 20133 Milano, Italy

²⁰Dipartimento di Astronomia, Università di Padova, vicolo dell'Osservatorio 2, I-35122 Padua, Italy

²¹Department of Astronomy, Columbia University, MC2457, 550 W. 120 St. New York, NY 10027

²²European Southern Observatory, Karl-Schwarzschild-Str. 2, D-85748 Garching, Germany

served colors (Baum 1962). This method is extremely efficient for assembling large redshift samples for faint galaxies. Despite having a lower accuracy than spectroscopic redshifts (hereafter, spectro-z), photo-z have the advantage of significantly improved completeness down to a flux limit fainter than the spectroscopic limit. Deep photo-z samples such as COMBO-17 (Wolf et al. 2003), CFHTLS (Ilbert et al. 2006), SWIRE (Rowan-Robinson et al. 2008), and COSMOS (Mobasher et al. 2007) contain more than 1,000,000 galaxies and go as faint as $i \sim 25$ with a relatively small amount of telescope time.

Typical photo-z with an accuracy of $\sigma_{\Delta z}/(1+z_s) \sim 0.02-0.04$ ($\Delta z = z_s - z_p$) are widely used to study the evolution of galaxy stellar masses and luminosities (e.g. Fontana et al. 2000; Wolf et al. 2003; Gabasch et al. 2004; Caputi et al. 2006; Arnouts et al. 2007), for angular clustering analysis (e.g. Heinis et al. 2007; McCracken et al. 2007), to study the relation between galaxy properties and environment (e.g. Capak et al. 2007), to trace large-scale structures (Mazure et al. 2007; Scoville et al. 2007) and to identify clusters at high redshift (Wang & Steinhardt 1998). Photo-z are also necessary for dark energy and dark matter weak lensing studies to separate foreground and background galaxies and to control systematic effects such as intrinsic shape alignment, shear shape correlation, and the effects of source clustering (Peacock et al. 2006). All of these applications require strict control of systematic effects in the photo-z estimate. An efficient way of identifying and removing systematics is to calibrate photo-z on a spectroscopic sample. The most common methods of calibration are neural network methods (e.g. Ball et al. 2004, Colister et al. 2004), a calibration of the color-z relation (Brodwin et al. 2006; Ilbert et al. 2006), or a reconstruction of the SED templates (Budavári et al. 2000; Feldmann et al. 2006).

As is also true for spectroscopic redshift measurements, photo-z accuracy depends on spectral coverage and resolution. It is also degraded for sources with a low signal-to-noise ratio (Bolzonella et al. 2000). The Balmer and Lyman breaks contain much of the photo-z information, so accuracy is lower in redshift ranges where these features are not well sampled by the filter set. As the photometric accuracy degrades, it becomes more

difficult to constrain the positions of these features, leading to lower accuracy. This creates a dual dependency of photo- z accuracy on magnitude and redshift which is defined by the survey design (exposure time, filter choice, and calibration accuracy). For this reason, any photo- z sample should be extensively tested and characterized in the same way a spectroscopic sample would be. This analysis is necessary in order to identify the redshift/magnitude ranges over which the photo- z can be trusted and used for a given scientific application.

This paper presents a new version of the photometric redshifts for the Cosmic Evolution Survey (COSMOS: Scoville et al. 2007) and an analysis of their accuracy. COSMOS is the largest *Hubble Space Telescope* (*HST*) survey ever undertaken - imaging an equatorial 2-deg² field to a depth of $I_{F814W} = 27.8$ mag (5σ , AB). The COSMOS field is equatorial to ensure visibility by all ground-based astronomical facilities. This project includes extensive multi-wavelength imaging from X-ray to radio (*XMM*, *Chandra*, *GALEX*, Subaru, CFHT, UKIRT, *Spitzer*, VLA). New ground-based NIR data (McCracken et al. 2008; Capak et al. 2008), *Spitzer*-IRAC data (Sanders et al. 2007), and medium/narrow band data from the Subaru Telescope (Taniguchi et al. 2007; Sasaki et al. 2008; Taniguchi et al. 2008; Capak et al. 2008) greatly improve the previous photometry catalogue (Capak et al. 2007). These new data are used for the photo- z derived here, yielding a factor of 3 higher accuracy than the first release of COSMOS photo- z (Mobasher et al. 2007).

The COSMOS data are presented in §2. The technique used to estimate the photo- z is presented in §3. In §4, we quantify the photo- z accuracy as a function of the magnitude and redshift. In §5, we provide the photo- z distribution of the i^+ selected samples. Specialized photo- z for X-ray selected sources, AGN, and variable objects, are discussed in a companion paper (Salvato et al. 2008).

Throughout this paper, we use the standard WMAP cosmology ($\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$) with $h = 0.7$ and $h = H_0/100$ km s⁻¹ Mpc⁻¹. Magnitudes are given in the AB system.

2. Data

Compared to the previous optical/near-infrared catalogue (Capak et al. 2007) the new photometry implements 14 new medium/narrow band data from the Subaru Telescope, deep ground-based NIR data (J and K bands) and *Spitzer*-IRAC data. The spectroscopic sample used to calibrate/test the photo- z is 10 times larger at $i_{AB}^+ < 22.5$ than that of Mobasher et al. (2007). The spectroscopic sample is supplemented with faint infrared selected sources and a deep faint spectroscopic sample at $z > 1.5$. Hereafter, we detail the photometric and spectroscopic data used to measure the photo- z .

2.1. Photometric data

Fluxes are measured in 30 bands from data taken on the Subaru (4200-9000Å), CFHT (3900-21500Å), UKIRT (12500Å), *Spitzer* (3.6-8 μ m) and *GALEX* (1500-2300Å) telescopes. We refer to Capak et al. (2008) for a complete description of the observations, data reduction and the photometry catalogue. The equivalent width and the effective wavelength of each filter are listed in Table 1. The sensitivities are given in Capak et al. (2008) and in Table 1 of Salvato et al. (2008). Table 1 indicates the fraction of sources detected in each band with an error less than 0.2 mag for a selection at $i^+ < 24.5$, $i^+ < 25$ and $i^+ < 25.5$. We summarize below the datasets used in this paper.

Ultraviolet: Very deep u^* band data were obtained at the 3.6m Canada-France Hawaii Telescope (CFHT) using the Megacam camera (Boulade et al. 2003). The u^* band data were processed at the TERAPIX data reduction center¹. The u^* band data cover the entire COSMOS field and reach a depth of $u^* \sim 26.5$ mag for a point source detected at 5σ . The u^* band images are also used as priors in the measurement of FUV (1500Å) and NUV (2300Å) fluxes in order to ensure a proper deblending of sources in the GALEX images (Zamojski et al. 2007). GALEX fluxes are then extracted using the EM-algorithm (Guillaume et al. 2006). They reach a depth of $FUV \sim 26$ mag and $NUV \sim 25.7$ mag.

Optical: The COSMOS-20 survey (Taniguchi et al. 2008) entailed 30 nights of observation at

¹terapix.iap.fr

the Subaru 8.2m telescope using the Suprime-Cam instrument. The observations are complete in 20 bands: 6 broad bands (B_J , V_J , g^+ , r^+ , i^+ , z^+), 12 medium bands ($IA427$, $IA464$, $IA484$, $IA505$, $IA527$, $IA574$, $IA624$, $IA679$, $IA709$, $IA738$, $IA767$, $IA827$) and 2 narrow bands ($NB711$, $NB816$).

Near-infrared: The catalogue includes deep J and K band data obtained using the WFCAM and WIRCAM wide-field infrared cameras on UKIRT and CFHT, respectively. The NIR data reduction is detailed in Capak et al. (2008) and McCracken et al. (2008). The data reach $J \sim 23.7$ mag and $K \sim 23.7$ mag for a point source detected at 5σ .

Mid-infrared: Deep IRAC data were taken during the *Spitzer* cycle 2 S-COSMOS survey (Sanders et al. 2007). A total of 166 hr were dedicated to cover the full 2-deg² with the IRAC camera in 4 bands: $3.6\mu\text{m}$, $4.5\mu\text{m}$, $5.6\mu\text{m}$ and $8.0\mu\text{m}$. Source detection is based on the $3.6\mu\text{m}$ image and the fluxes were measured in the four IRAC bands using the “dual mode” configuration of SExtractor. The IRAC catalogue is 50% complete at $1\mu\text{Jy}$ at $3.6\mu\text{m}$ ($m_{3.6\mu\text{m}} \sim 23.9$ mag).

All of the imaging data were combined to generate a master photometry catalogue (Capak et al. 2008). Photometry was done using SExtractor in dual mode (Bertin & Arnouts 1996). Source detection was run on the deepest image ($i^+ \sim 26.2$ for a point source detected at 5σ). For the UV-NIR data, the Point Spread Function (PSF) varies from $0.5''$ to $1.5''$ from the K to the u^* images. In order to obtain accurate colors, all the images were degraded to the same PSF of $1.5''$ following the method described in Capak et al. (2007). The final photometry catalogue contains PSF matched photometry for all the bands from the u^* to the K band, measured over an aperture of $3''$ diameter at the position of the i^+ band detection. For the FUV and NUV data, we transformed the total flux provided for the *GALEX*-FUV and -NUV counterpart by multiplying it by a factor 0.759. This factor is the fraction of the flux observed in optical into a $3''$ aperture flux as determined for point sources from simulations by Capak et al. (2007). To compensate for this approximation, we add in quadrature 0.1 and 0.3 mag to the all the measured errors in the *GALEX* NUV and FUV bands.

For the MIR data we did not degrade the optical data to the larger IRAC PSF. Instead, the fol-

lowing procedures were used. An IRAC flux was first associated with the optical sources by matching their positions in each of the 4 IRAC bands within a search radius of $1''$. Following Surace et al. (2004), the IRAC fluxes were then measured in a circular aperture of radius $1.9''$. The aperture flux was then converted to a total flux using the aperture correction factors 0.76, 0.74, 0.62, 0.58 at $3.6\mu\text{m}$, $4.5\mu\text{m}$, $5.6\mu\text{m}$ and $8.0\mu\text{m}$, respectively (Surace et al. 2004). Since the optical fluxes were measured over an aperture of $3''$ diameter which encloses $\sim 75\%$ of the flux for a point like source, we then multiply the IRAC total fluxes by a factor 0.75. This approximation provides good agreement between the predicted and observed colors ($z^+ - 3.6\mu\text{m}$ and $3.6\mu\text{m} - 4.5\mu\text{m}$ colors). In order to compensate for this approximation, we add in quadrature 0.1, 0.1, 0.3 and 0.3 mag to the errors in the IRAC 3.6, 4.5, 5.6 and $8.0\mu\text{m}$.

Finally, all magnitudes are corrected for galactic extinction estimated for each object individually, using dust map images from Schlegel et al. (1998). We limit the photo- z analysis to an area of 2-deg² ($1.49878 < \alpha < 2.91276$ and $149.41140 < \delta < 150.826934$) which has a uniform and deep coverage in all the bands. Poor image quality areas (e.g. field boundary, saturated stars, satellite tracks and image defects) are masked. Photo- z are computed only in the non-masked regions with a total covered area of 1.73-deg². 126 071, 293 627 and 607 617 sources are detected at $i^+ < 24$, $i^+ < 25$ and $i^+ < 26$.

2.2. Spectroscopic data

The spectroscopic samples were observed with the VIMOS/VLT spectrograph (zCOSMOS; Lilly et al. 2007) and the DEIMOS/Keck spectrograph (Kartaltepe et al. 2008). These two spectroscopic samples have very different selection criteria (see below); they therefore cover very different ranges of redshift and color space, providing a broad sample for evaluation of the photo- z .

The zCOSMOS survey (Lilly et al. 2007) has two components: zCOSMOS-bright with a sample of 20,000 galaxies selected at $i^* \leq 22.5$ and zCOSMOS-faint with approximately 10,000 galaxies color-selected to lie in the redshift range $1.5 \lesssim z \lesssim 3$. In the latter, galaxies are selected by color based either on the BzK criterion (Daddi et al. 2004) or the *UGR* “BM” and “BX” crite-

filter	telescope	effective λ	FWHM	s_f	% at $i^+ < 24.5$	% at $i^+ < 25$	% at $i^+ < 25.5$
u^*	CFHT	3911.0	538.0	0.054	89.3	85.2	77.2
B_J	Subaru	4439.6	806.7	-0.242	97.1	95.2	90.5
V_J	Subaru	5448.9	934.8	-0.094	99.3	98.2	94.2
g^+	Subaru	4728.3	1162.9	0.024	96.4	93.6	86.0
r^+	Subaru	6231.8	1348.8	0.003	99.6	99.5	98.4
i^+	Subaru	7629.1	1489.4	0.019	99.9	99.9	99.8
i^*	CFHT	7628.9	1460.0	-0.007	37.8	25.4	17.4
z^+	Subaru	9021.6	955.3	-0.037	99.8	97.9	83.8
J	UKIRT	12444.1	1558.0	0.124	65.4	49.1	35.7
K_S	NOAO	21434.8	3115.0	0.022	15.3	10.3	7.08
K	CFHT	21480.2	3250.0	-0.051	84.1	68.5	52.1
$IA427$	Subaru	4256.3	206.5	0.037	77.1	64.3	48.4
$IA464$	Subaru	4633.3	218.0	0.013	78.5	64.3	47.6
$IA484$	Subaru	4845.9	228.5	0.000	88.7	78.3	62.0
$IA505$	Subaru	5060.7	230.5	-0.002	84.0	70.0	52.0
$IA527$	Subaru	5258.9	242.0	0.026	93.2	84.6	68.7
$IA574$	Subaru	5762.1	271.5	0.078	92.5	80.0	60.6
$IA624$	Subaru	6230.0	300.5	0.002	97.4	90.2	72.2
$IA679$	Subaru	6778.8	336.0	-0.181	99.5	96.7	82.9
$IA709$	Subaru	7070.7	315.5	-0.024	99.8	97.7	83.5
$IA738$	Subaru	7358.7	323.5	0.017	99.5	94.2	73.6
$IA767$	Subaru	7681.2	364.0	0.041	99.6	93.5	72.0
$IA827$	Subaru	8240.9	343.5	-0.019	99.7	97.2	81.8
$NB711$	Subaru	7119.6	72.5	0.014	84.8	60.7	41.9
$NB816$	Subaru	8149.0	119.5	0.068	99.8	99.1	88.2
$IRAC1$	<i>Spitzer</i>	35262.5	7412.0	0.002	70.6	60.8	48.4
$IRAC2$	<i>Spitzer</i>	44606.7	10113.0	0.000	62.6	51.6	39.7
$IRAC3$	<i>Spitzer</i>	56764.4	13499.0	0.013	33.7	26.0	19.5
$IRAC4$	<i>Spitzer</i>	77030.1	28397.0	-0.171	15.7	11.3	8.1
FUV	<i>GALEX</i>	1551.3	230.8	0.314	8.5	5.8	4.0
NUV	<i>GALEX</i>	2306.5	789.1	-0.022	19.7	13.4	9.2

Table 1: Effective wavelength, width and systematic offsets s_f in magnitude (with our definition, s_f have the opposite sign to Table.13 of Capak et al. 2007) plus the fraction of sources in each band with an error less than 0.2 magnitude at $i^+ < 24.5$, $i^+ < 25$, $i^+ < 25.5$.

rion of Steidel et al. (2004), and the magnitude cut was $B_J < 24 - 25$ (depending on the color cut). zCOSMOS-bright galaxies were observed using the red grism of VIMOS covering a wavelength range $5500\text{\AA} < \lambda < 9000\text{\AA}$ at a resolution of 600 (MR grism). For the zCOSMOS-faint sample, observations were carried out with the blue grism of VIMOS ($3600\text{\AA} < \lambda < 6800\text{\AA}$) at a resolution of 200.

The zCOSMOS-bright survey is now $\sim 50\%$ complete. Here we make use of only the extremely secure spectro-z measurements with a confidence level greater than 99% (class 3 and 4). This secure zCOSMOS-bright sample contains 4148 galaxies with a median redshift of ~ 0.48 . The zCOSMOS-faint survey is in its early stages, and here we use a preliminary sample of 148 galaxies with a median redshift of $z_m \sim 2.2$ and as faint as $i^+ \sim 25$. This zCOSMOS-faint spectroscopic sample is not fully representative of the average population at $1.5 < z < 3$ due to the selection criteria (e.g. $B_J < 24 - 25$).

The Keck II spectroscopic follow-up of $24\mu\text{m}$ selected sources (Kartaltepe et al. 2008) is on-going and we refer to this sample as MIPS-spectro-z. The DEIMOS spectra cover a wavelength range $4000\text{\AA} < \lambda < 9000\text{\AA}$ at a resolution of 600. This sample of $24\mu\text{m}$ selected galaxies contains 317 secure spectro-z (at least two spectral features) with an average redshift of $z \sim 0.74$ and apparent magnitude in the range $18 < i^+ < 25$.

For all of the spectroscopic samples used in this paper for testing and verification of the photo-z, we include only secure spectro-z. Therefore, the uncertainties in the spectro-z are neglected and the spectro-z are used as a reference to assess the quality of the photo-z.

3. Photometric Redshift derivation

Photometric redshifts were derived using the *Le Phare* code² (S. Arnouts & O. Ilbert) which is based on a χ^2 template-fitting procedure. In the discussion below, we focus on the improvements introduced here as compared to Ilbert et al. (2006) and the previous COSMOS photo-z (Mobasher et al. 2007).

²www.oamp.fr/people/arnouts/LE_PHARE.html

3.1. Galaxy SED template library




Fig. 1.— SED templates. The flux scale is arbitrary. The top 12 SEDs (cyan) are generated with Bruzual & Charlot (2003). The spiral (green) and elliptical (red) SEDs are from Polletta et al. (2007).

Ilbert et al. (2006) and Mobasher et al. (2007) used a set of local galaxy SED templates (CWW: Coleman, Wu, & Weedman 1980) which have been widely employed for photo-z (e.g. Sawicki 1997; Fernández-Soto et al. 1999; Arnouts et al. 1999; Brodwin et al. 2006). Here, we employ a new set of templates generated by Polletta et al. (2007) with the code GRASIL (Silva et al. 1998). Polletta et al. selected their templates for fitting the VVDS sources (Le Fèvre et al. 2005) from the UV-optical (CFHTLS: McCracken et al. 2007) to the mid-IR (SWIRE: Lonsdale et al. 2003). Therefore, this set of templates provides a better joining of UV and MIR than those by CWW. The 9 galaxy templates of Polletta et al. (2007) include 3 SEDs of elliptical galaxies and 6 templates of spiral galaxies (S0, Sa, Sb, Sc, Sd, Sdm).

We did find that the blue observed colors of the spectroscopic sample were not fully reproduced by the Polletta et al. (2007) templates. We therefore generated 12 additional templates using Bruzual & Charlot (2003) models with starburst (SB) ages

ranging from 3 Gyr to 0.03 Gyr. We extend the BC03 templates beyond $3\mu\text{m}$ rest-frame using the Sdm template of Polletta et al. (2007). The full library of template SEDs, 9 from Polletta et al. (2007) and 12 from Bruzual & Charlot (2003), is shown in Fig. 1. Finally, we linearly interpolated between some Polletta et al. templates to refine the sampling in $color - z$ space.

Figure 2 shows the observed colors and redshifts of the spectroscopic sample compared with the predicted colors for the library SEDs.

3.2. Emission lines

Fig. 2 clearly shows how the observed colors oscillate with the redshift, especially when the colors are measured with the medium bands (top panels). Comparing the template curves with (dashed) and without (solid) emission lines, one sees that the expected line fluxes can cause up to 0.4 mag changes in the color. This effect is particularly important when colors involving intermediate and narrow band filters are computed (see for example the upper-right panel in Fig. 2) but can be already noticed using broad band colors. The color oscillations are well explained by the contribution of emission lines like $H\alpha$, $[O\ III]$, and $[O\ II]$ to the observed flux; thus the contribution of the emission lines to the flux must be taken into account to obtain accurate photo- z and this is a major change implemented here compared to Ilbert et al. (2006) and Mobasher et al. (2007).

In order to include the emission line contribution to the SED, we need to model the emission line fluxes ($O\ II$, $O\ III$, $H\beta$, $H\alpha$, $Ly\alpha$) at any redshift, template and extinction. The rescaling of the template (A in eqn.2) determines also the emission line fluxes (therefore, the modeling of the fluxes must be done galaxy per galaxy).

Our new procedure estimates the $[O\ II]$ emission line flux from the UV luminosity of the rescaled template, using the Kennicutt (1998) calibration laws. In the template fitting, a UV rest-frame luminosity corrected for dust extinction can be computed at every step of the redshift/template/extinction grid (the rescaling factor A is taken into account in the UV luminosity). The UV luminosity (at $2300\text{\AA}$) is then related to the SFR using the relation $SFR(M_\odot\ yr^{-1}) = 1.4 \times 10^{-28} L_\nu(erg\ s^{-1} Hz^{-1})$

from Kennicutt (1998). This SFR can then be translated to an $[O\ II]$ emission line flux using the relation $SFR(M_\odot\ yr^{-1}) = (1.4 \pm 0.4) \times 10^{-41} L_{[OII]}(erg\ s^{-1})$ (Kennicutt 1998). This translates to :

$$\log(F_{[OII]}) = -0.4 \times M_{UV} + 10.65 - \frac{DM(z)}{2.5} \quad (1)$$

where DM is the distance modulus, $F_{[OII]}$ is expressed in units of $10^{-17} erg\ s^{-1} cm^2$ and M_{UV} is the dust corrected $UV(2300\text{\AA})$ absolute magnitude.




Fig. 3.— Relation between the $[O\ II]$ flux and the absolute magnitudes in UV ($2300\text{\AA}$). The solid line corresponds to the relation obtained by applying the Kennicutt (1998) relations between SFR_{OII} and SFR_{UV} as used here to include emission line fluxes in the photo- z fitting (see eqn.1) and the points are observed emission line fluxes from VVDS (Lamareille et al. 2008). (The UV luminosities and $O\ II$ fluxes are corrected for dust extinction.) The red and blue points are galaxies with $M_U - M_R > 1.6$ and $M_U - M_R < 1.6$, respectively. The larger symbols correspond to larger equivalent width.

Figure 3 shows the measured $[O\ II]$ fluxes from VVDS (Lamareille et al. 2008) and the relation (solid line) expected from Kennicutt (1998). We can perform this comparison only at $0.5 < z < 1.4$,




Fig. 2.— The observed colors and redshifts (z_s) for the spectroscopic sample galaxies (open stars). The solid lines are the predicted colors as a function of redshift for some SEDs of the library (red: Elliptical and green: Spiral from Polletta et al. 2007 ; cyan: Bruzual & Charlot 2003). The solid curves are the predicted colors without including emission lines (no reddening for the elliptical templates, $E(B - V) = 0.2$ for the late types) whereas the dashed curves are the same templates including the emission line fluxes (assuming $M_{FUV} = -20$ in this example). The top right panel clearly shows that the emission lines can change the colors up to 0.4 mag.

when the [O II] line is observable in the VIMOS spectra. Figure 3 shows the good correlation between the measured and predicted [O II] fluxes, with an rms dispersion of 0.2 dex. In acknowledgment of this dispersion, we allow the intrinsic [O II] flux to vary by a factor of 2 in the templates with respect to the nominal flux predicted by eqn.1.

With our procedure, we can predict for each galaxy the [O II] flux at every redshift, template and extinction combination. For the other emission lines, we adopt intrinsic, unextincted flux ratios of [O III/O II] = 0.36; [H β /O II] = 0.61; [H α /O II] = 1.77 and [Ly α /O II] = 2 (McCall et al. 1985, Moustakas et al. 2006, Mouhcine et al. 2005, Kennicutt 1998). When we apply an additional extinction to the template, we modify these ratios with the corresponding attenuation. Then, we sum the emission line fluxes to the template continuum before integrating through the filter transmission curves.

The effect of these emission lines on the modeled *color-redshift* relation is shown in Fig. 2 for a galaxy at $M_{UV} = -20$. The oscillations in the observed colors versus redshift are well reproduced by the models.

3.3. Systematic offsets

The χ^2 template-fitting method is meaningful only if the *color-z* relation predicted from the templates is a good representation of the observed *color-z* relation. Uncertainties in the zero-point offsets of photometric bands can lead to systematic shifts between the predicted *color-z* relation and the observed colors of the spectroscopic sample.

To evaluate the zero-point errors, we use the spectroscopic sample and set the redshift to the spectro-z value. Then, we determine the best-fit template for each spectroscopic galaxy. For random, normally distributed uncertainties in the flux measurements, $\overline{\Delta F_f}$ should be ~ 0 (where $\overline{\Delta F_f}$ is the average difference between predicted and observed fluxes in the filter f). Instead, we initially find systematic offsets of $\overline{\Delta F_f}$ (as for earlier photo-z derivations, e.g. Brodwin et al. 2006; Ilbert et al. 2006; Mobasher et al. 2007). Such offsets are mainly due to:

- uncertainties in the absolute calibration of the photometric zero-points.

- uncertainties in the color modeling (filter transmission curves, incomplete set of templates or an incorrect extinction curve).

These systematic offsets were removed using the iterative procedure detailed in Ilbert et al. (2006). For each filter, f , we estimate the values s_f which minimize $\overline{\Delta F_f}$. After having applied the corrections, s_f , in each band f , the systematic offsets derived in a second iteration can change. The values converge after 3 iterations and we adopted the systematic offsets listed in Table. 1.

3.4. Extinction law

Ilbert et al. (2006) adopted the dust extinction law measured in the Small Magellanic Cloud (Prevot et al. 1984). However, considerable changes in the extinction curve are expected from galaxy to galaxy. Maraston et al. (2006) considered as a free parameter the different extinction curves of the “Hyperz” package (Bolzonella et al. 2000) (e.g. Milky Way, large and small Magellanic clouds and Calzetti). In order to limit the risk of catastrophic failures, we adopt an intermediate approach here, using the most suitable extinction curve depending on the SED template.

For each galaxy of the zCOSMOS sample, we set the redshift to the spectroscopic redshift value. Then, we determine the best fit-template and the appropriate color excess $E(B-V)^{\text{best}}$. In this fit, we assume an extinction curve $k(\lambda)$ ($= A(\lambda)/E(B-V)$). Since the extinction curves do not differ strongly at $\lambda > 3000\text{\AA}$ (blue and green curves in Fig. 4), we fit the templates using only passbands with $\lambda > 3000(1+z)\text{\AA}$. With this procedure, the $E(B-V)^{\text{best}}$ value does not depend significantly on the adopted extinction curve.

The extinction curves differ strongly at $\lambda < 3000\text{\AA}$. Therefore, we use the rest-frame observed SEDs at $\lambda_{\text{rest-frame}} < 3000\text{\AA}$ to discriminate between the different extinction curves. Using m^{obs} to represent the observed magnitude and $m_{\text{uncor}}^{\text{template}}$ to represent the predicted magnitude from the best-fit template (uncorrected for extinction), the extinction $A(\lambda)$ is given by $A(\lambda) = m^{\text{obs}} - m_{\text{uncor}}^{\text{template}}$. Fig. 4 shows the rest-frame $(m^{\text{obs}} - m_{\text{uncor}}^{\text{template}})/E(B-V)^{\text{best}}$ for each flux measurement, i.e the extinction curve $k(\lambda)$. These points are compared with the Calzetti et al. (2000) and the Prevot et al. (1984) extinction curves.




Fig. 4.— Attenuation by dust as a function of λ . The points are the extinction $A_i/(E_i(B - V))$ estimated from the galaxies with a spectro- z (see §3.4). The red points at $\lambda < 3000\text{\AA}$ are not used to estimate $E_i(B - V)$. The Prevot et al. (1984) and Calzetti et al. (2000) extinction curves are shown with the blue and green solid lines, respectively. The Prevot et al. (1984) extinction law is rescaled to the same A_V as the Calzetti law by applying a factor $4.05/2.72$. The extinction curve derived by Prevot et al. (1984) is used for the galaxies redder than SB3 and the Calzetti et al. (2000) extinction law for the galaxies bluer than SB3.

(We scaled the Prevot et al. curve to the same A_V value as Calzetti et al. by applying a factor $4.05/2.72$ which is the ratio between the R_V of the Calzetti and Prevot laws.) The Small Magellanic Cloud extinction curve (Prevot et al. 1994) is well suited for galaxies redder than the starburst template SB3. For galaxies bluer than SB3, the Calzetti et al. (2000) extinction curve is found to be more appropriate (see Fig. 4). For the IR sources which are strongly star-forming, Caputi et al. (2008) show also that the Calzetti et al. (2000) extinction law is more appropriate. This result is not surprising since the Calzetti law was determined from observed starburst galaxies. A broad absorption excess at $2175\text{\AA}$ (UV bump) seems necessary to explain the UV flux in some starburst galaxies. The presence of this UV bump can be seen in a theoretical modeling of the Calzetti law (Fischera et al. 2004) and in the K20 sample of high redshift galaxies at $1 < z < 2.5$ (Noll et al. 2007).

To summarize, we apply an additional extinction to the templates according to a grid $E(B - V) = 0, 0.05, 0.1, 0.15, 0.2, 0.25, 0.3, 0.4, 0.5$. We use the Prevot et al. (1994) extinction curve for the templates redder than SB3, and Calzetti et al. (2000) for the templates bluer than SB3. We allow an additional bump at $2175\text{\AA}$ for the Calzetti extinction law if it produces a smaller χ^2 . No reddening is allowed for galaxies redder than *Sb*.

3.5. χ^2 minimization

The photo- z is the redshift value which minimizes the merit function $\chi^2(z, T, A)$:

$$\chi^2 = \sum_{f=1}^{N_f} \left(\frac{F_{\text{obs}}^f - A \times F_{\text{pred}}^f(z, T) 10^{-0.4s_f}}{\sigma_{\text{obs}}^f} \right)^2, \quad (2)$$

where $F_{\text{pred}}^f(T, z)$ is the flux predicted for a template T at redshift z . F_{obs}^f is the observed flux and σ_{obs}^f is the associated error. The index f refers to each specific filter and s_f is the zero-point offset listed in Table.1. The opacity of the inter-galactic medium (Madau et al. 1995) is taken into account. The photo- z is estimated from the minimization of χ^2 with respect to the free parameters, z , T and the normalization factor A . The color excess $E(B - V)$ is included in the term T (see section 3.4). The grid spacing in redshift is $\delta z = 0.01$ and

the final redshift is derived by parabolic interpolation of the redshift probability distribution. The redshift probability distribution function (PDF z) is derived directly from the $\chi^2(z)$ distribution:

$$P(z) \propto \exp\left(-\frac{\chi^2(z) - \chi_{\text{min}}^2}{2}\right). \quad (3)$$

The minimum and maximum redshifts around the photo- z solution, corresponding to the 1σ errors, are estimated from the equation $\chi^2(z) = \chi_{\text{min}}^2 + 1$. As in Ilbert et al. (2006), we increased the SExtractor flux errors by a factor of 1.5. This factor does not shift the best-fit photo- z value but broadens the χ^2 peak and derived redshift uncertainty.




Fig. 5.— Color-color diagnostics for sources including stars and galaxies. Stars revealed by the χ^2 classification are shown in green; the red and grey points are galaxies with $z_p > 1$ and $z_p < 1$, respectively.

3.6. Star/AGN/galaxy classification

For each object, χ^2 is evaluated for both the galaxy templates and stellar SED templates (Chabrier et al. 2000, Bixler et al. 1991). If $\chi_{\text{gal}}^2 - \chi_{\text{star}}^2 > 0$, where χ_{gal}^2 and χ_{star}^2 are the minimum χ^2 values obtained with the galaxy and stellar templates respectively, the object is flagged as a possible star. Leauthaud et al. (2007) catalogued point-like sources in the COSMOS field

using the peak surface brightness measured on the ACS images. The SED χ^2 and morphological classification methods were found to be in excellent agreement – 84% of the point-like sources at $i^+ < 24$ are classified as stars with the SED χ^2 criterion mentioned above, while only 0.2% of the extended sources on the ACS images are misclassified as stars with the same χ^2 criterion. As shown in Fig. 5, the star sequence colors are well distinguished from the bulk of the galaxy population if the colors include a NIR band. NIR and MIR data are crucial to separate stars from galaxies. Therefore, we limit the χ^2 star classification to $K < 24$ or $F_{3.6\mu m} > 1\mu Jy$. 2% of the point-like sources with $i^+ < 24$ have $K > 24$ and $F_{3.6\mu m} > 1\mu Jy$. These objects, which are only 2% of the total population at $i^+ < 24$, could be stars not recognized as such by our SED χ^2 classification.

AGN can be identified by their point-like X-ray emission. Most ($\sim 90\%$) of the 1887 sources detected in *XMM*-COSMOS (Brusa et al. 2007) are dominated by an AGN. Their photo-z determinations require a different treatment: a correction for variability of the photometric data and the use of SED templates specifically tuned to AGN and their host galaxies. Accurate photo-z for the *XMM*-COSMOS sources are derived in a companion paper (Salvato et al. 2008). Due to the flux limit in *XMM*-COSMOS (Brusa et al. 2007), the moderately luminous ($\log(L_x)=42-43$ erg/s) and luminous ($\log(L_x)=43-44$ erg/s) AGN are not sampled by *XMM*-COSMOS at $z = 0.5 - 1.25$ and $z = 1.5 - 3$, respectively. Therefore, this population is not identified as AGN in the galaxy catalogue.

4. Photometric redshift accuracy

In this section, we assess the quality of the derived photometric redshifts by two approaches: comparison with high confidence spectroscopic redshifts and an analysis of the width of the redshift probability distribution function obtained from the χ^2 fitting for the photo-z. The latter approach is justified by the excellent agreement of the z_s - z_p distribution with the width of the probability distribution function for the spectroscopic sample.

4.1. Comparison of Photometric and Spectroscopic Redshifts

We first assess the quality of the photo-z by comparison with the spectro-z. If $\Delta z = z_s - z_p$, we can estimate the redshift accuracy from $\sigma_{\Delta z/(1+z_s)}$ using the normalized median absolute deviation (NMAD; Hoaglin et al. 1983) defined as $1.48 \times \text{median}(|z_p - z_s|/(1+z_s))$. The NMAD is directly comparable to other papers which directly quote the $rms/(1+z)$. This dispersion estimate is robust with respect to catastrophic errors (i.e., objects with $|z_p - z_s|/(1+z_s) > 0.15$). The percentage of catastrophic errors is denoted by η .

Figure 6 (left panel) shows the comparison between z_p and z_s for the zCOSMOS-bright sample selected at $i^+ < 22.5$. The spectro-z sample is selected only by apparent magnitude and is therefore representative of the entire $i^+ < 22.5$ population. We obtain an accuracy of $\sigma_{\Delta z/(1+z)} = 0.007$ at $i^+ < 22.5$ and the distribution of offsets $((z_p - z_s)/(1+z_s))$ is well fit by a gaussian with $\sigma = 0.007$ (Fig. 6 right panel). The percentage of catastrophic failures is below 1%.

The zCOSMOS-faint sample (Fig. 7, left panel) with a median apparent magnitude of $i_{\text{med}}^+ \sim 24$ provides a quality check for the photo-z at $1.5 < z < 3$ where the photo-z are expected to have a significantly higher uncertainty. The faint sample includes galaxies at $i^+ \sim 25$ and the Balmer break is shifted into the NIR where the filter set has gaps. At $1.5 < z < 3$, the accuracy is found to be $\sigma_{\Delta z/(1+z)} = 0.06$ with 20% catastrophic failures. The fraction of high-redshift galaxies ($z_s > 1.5$) for which a low-redshift photo-z ($z_p < 0.5$) was assigned is 7% but this failure rate drops to 4% if the sample is restricted to galaxies detected at $IRAC(3.6\mu m) > 1\mu Jy$. The zCOSMOS-faint sample actually includes 54 galaxies with low spectroscopic redshift ($z_s < 0.5$); 53 out of the 54 galaxies ($> 98\%$) were assigned the correct photo-z low redshift. In summary, we conclude that, for the faint galaxies, the photo-z of low redshift objects are still assigned correctly, while the failure rate for high redshift objects is significantly reduced if the objects have good IRAC detections. Such result is expected since the photo-z are already including the same information present in high-z color selections such as *BzK* (Daddi et al. 2004) or similar diagnostics using the IRAC bands




Fig. 6.— Left panel: Comparison between z_p and z_s for the bright spectroscopic selected sample $17.5 \leq i_{AB}^+ \leq 22.5$ (zCOSMOS-bright: Lilly et al. 2008). The dotted and dashed lines are for $z_p = z_s \pm 0.15(1 + z_s)$ and $z_p = z_s \pm 0.05(1 + z_s)$, respectively. The 1σ dispersion, the fraction of catastrophic failures and the median apparent magnitude are listed in the top left corner of the left panel. Right panel: $\Delta z / (1 + z)$ distribution. The dashed line is a gaussian distribution with $\sigma = 0.007$.




Fig. 7.— Comparison between z_p and z_s . Left panel: zCOSMOS-faint sample (Lilly et al. 2008). The open triangles are objects with a secondary peak in the redshift probability distribution function. Right panel: infrared selected sample (Kartaltepe et al. 2008) split into a bright sample $i^+ < 22.5$ (black), a faint sample $22.5 < i^+ < 24$ (red) and a very faint sample $24 < i^+ < 25$ (green).

(Fig. 5) to isolate the good redshift range.

The zCOSMOS-bright and zCOSMOS-faint samples do not probe $0.2 < z < 1.5$ at $i^+ > 22.5$. Here, we use as a comparison sample the MIPS-spectro-z (Kartaltepe et al. 2008). In Fig. 7 (right panel), we split the MIPS-spectro-z sample into bright ($i^+ < 22.5$), faint ($22.5 < i^+ < 24$) and very faint samples ($24 < i^+ < 25$). For the bright sub-sample, the dispersion of the MIPS-selected galaxies is $\sigma_{\Delta z/(1+z_s)} = 0.009$, only slightly greater than that of the optically selected sample at $i^+ < 22.5$. For the faint sub-sample (median apparent magnitude $i^+ \sim 23.1$), $\sigma_{\Delta z/(1+z_s)} = 0.011$, i.e. slightly worse than for the brighter optically-selected objects. For the very faint sub-sample, the accuracy is degraded to $\sigma_{\Delta z/(1+z_s)} = 0.053$ with a catastrophic failures rate of 20%. This degradation is due to decreasing signal-to-noise photometry for faint objects and could be amplified by the infrared selection which picks up more heavily obscured galaxies and higher redshift galaxies (e.g. Fig.4 of Le Floch et al. 2005).




Fig. 8.— Cumulative distribution of the ratio $|z_p - z_s|/(1\sigma \text{ error})$. 65% of the photo-z have a spectro-z solution encompassed within the 1σ error, close to the expected value of 68% (magenta dashed line).




Fig. 9.— 1σ uncertainty for the z_p estimate as a function of redshift in different apparent magnitude bin. Each value is computed with 50 galaxies per bin. The thick solid green lines, the solid red line, the dashed blue lines and the dotted magenta lines are for $i^+ < 22.5$, $22.5 < i^+ < 24$, $24 < i^+ < 25$, $25 < i^+ < 25.5$, respectively.

4.2. Accuracy derived from the Photo-z Probability Distribution Function

Since evaluation of the photo-z accuracy from the comparison with spec-z is limited to specific ranges of magnitude and redshift, we use the 1σ uncertainty in the derived photo-z probability distribution to extend the uncertainty estimates over the full magnitude/redshift space.

The reliability of the 1σ uncertainty estimate for the photo-z as derived from the Probability Distribution Function (PDFz) (see §3.5) can be checked by comparing this uncertainty with that derived directly from the photo-z - spec-z offsets for the spectroscopic sample. Figure 8 shows the cumulative distribution of these offsets normalized by the 1σ uncertainty in the probability function for the zCOSMOS-bright sample (the ratio $|z_p - z_s|/(1\sigma \text{ error})$ is lower than 1 if the measured offset $z_p - z_s$ is lower than the 1σ uncertainty). 65% of the z_p are within the 1σ error bars, whereas the expected fraction is 68%. We

therefore conclude that the 1σ uncertainties in the probability function as derived here provide a robust assessment of the accuracy in z_p .

Figure 9 shows the 1σ negative and positive uncertainties derived from the probability function as a function of redshift and apparent magnitude. Two clear conclusions emerge: the accuracy is inevitably degraded for fainter galaxies at all redshifts and the photo- z have significantly higher uncertainty at $z \gtrsim 1.25$.

From $z > 0$ out to $z = 1.25$, the 1σ errors do not depend significantly on the redshift, with $\sigma_{\Delta z} \lesssim 0.02$ at $i^+ < 24$ (note that we dropped out the division of σ by $(1+z)$ in this analysis). To first order, the photo- z are accurate when the wavelengths of the Balmer and/or Lyman breaks are well constrained. Therefore, the wavelength coverage of the filter set and the photometry sensitivities determine the photo- z accuracy as a function of redshift. The COSMOS photometry coverage is continuous and dense in the optical from the u^* band ($\lambda_{\text{eff}} \sim 3911\text{\AA}$) to the z^+ band ($\lambda_{\text{eff}} \sim 9021\text{\AA}$). The average wavelength spacing between consecutive filters is also only $230\text{\AA}$. The Balmer break at redshift $z = 0$ out to $z = 1.25$ is always at $\lambda < 9000\text{\AA}$, thus the high accuracy in this redshift range.

The accuracy degrades at $z > 1.25$, where the $4000\text{\AA}$ Balmer break goes out of the z^+ band; at $z \sim 1.8$ it is a factor 3 larger than at $z \sim 1.25$ ($\sigma_{\Delta z} \sim 0.14$ for $i^+ \sim 24$). The lack of coverage between the z' and the J band accounts for the discontinuous increase in uncertainty at $z > 1.25$ since the Balmer break can not be located with precision. At $z > 1.5$, the estimated error from the photo- z – spectro- z comparison is $\sigma_{\Delta z} = 0.19$ (see §4.1; $\sigma_{\Delta z/(1+z_s)} = 0.06$ at the median redshift $z \sim 2.2$). This accuracy is ~ 1.4 bigger than that estimated from the 1σ uncertainty ($\sigma_{\Delta z} \sim 0.14$ at $i^+ \sim 24$). Errors due to bias in photo- z will not be recovered from the probability distribution function which could explain this difference. At $z > 1.25$, all the optical filters, which are about 80% of all used filters, are sampling the rest-frame UV at $\lambda < 3500\text{\AA}$. Uncertainties in the adopted extinction law have considerable impact on the UV slope and could introduce small biases in the estimate of the photo- z .

At $z > 2.5$, the accuracy improves improves again ($\sigma_{\Delta z} \lesssim 0.1$ for $24 < i^+ < 25$) when the

$4000\text{\AA}$ Balmer break enters in the J band ($z \sim 2$) and the UV light shortward of L_α enters the u^* band ($z \sim 2.3$).

4.3. Importance of zero-point offsets and emission lines

The major improvements in the technique implemented here are the carefully iterative evaluation of the photometric zero-point offsets for all bands and the allowance for a range of emission line contributions to the fluxes of template SEDs.

The left panel of Fig. 10 shows the comparison between z_p and z_s computed without correction of the systematic band offsets (see §3.3) and without including the emission lines (see §3.2). Some systematic biases (horizontal and vertical stripes in Fig. 10) greater than $\delta_z \sim 0.1$ are clearly seen in the photo- z estimate. For example, galaxies with $z_s \sim 0.8$ are often shifted to $z_z \sim 0.7$. With the iterative calibration of the band offsets turned on (right panel of Fig. 10), these biases are limited to $\delta_z < 0.05$. This clearly demonstrates the importance of zero-point calibration to reduce the photo- z bias (as already shown by Brodwin et al. 2006 and Ilbert et al. 2006).

The right panel of Fig. 10 shows the comparison between z_p and z_s without including the emission lines (see §3.2). The accuracy is $\sigma_{\Delta z/(1+z_s)} \sim 0.02$. When the emission lines are included in the templates, the accuracy is improved by a factor of 2.5 ($\sigma_{\Delta z/(1+z_s)} = 0.007$, left panel of Fig. 6). Therefore, including emission lines in the SEDs is crucial, especially when the medium bands are used to measure z_p .

5. Discussion

5.1. Comparison with Other Surveys

We now compare the accuracy of the COSMOS-30 photo- z (this work) with those obtained previously for COMBO-17 (Wolf et al. 2004) and CFHTLS-DEEP (Ilbert et al. 2006). The accuracies can be compared as a function of apparent magnitude using the 1σ measured uncertainties in the photo- z probabilities. Once again, we check also the validity of the 1σ uncertainties for other surveys following the method described in §4.2. For the COMBO-17 and CFHTLS photo- z , we use the spectro- z from the VIMOS-VLT Deep Survey




Fig. 10.— Comparison between z_p and z_s at different steps of the method. The comparison is done for the zCOSMOS-bright sample selected at $17.5 \leq i^+ \leq 22.5$. The left panel shows the photo- z computed with a standard χ^2 method (no emission lines contributions and without calibration of the band offsets). The right panel shows z_p computed with the calibration of the band offsets, but without including emission lines in the templates. Including emission lines improves the accuracy by a factor ~ 2.5 .

(Le Fèvre et al. 2004, 2005). Figure 11 shows that about 53%, 67% of the values for z_s are included inside the 1σ uncertainties for COMBO-17 and CFHTLS-DEEP survey, respectively. The 1σ errors for z_p in COMBO-17 are rescaled by ~ 1.2 to obtain 68% of the z_s within the 1σ error (This rescale is small and changes nothing in our conclusions).

Figure 12 shows the redshift dependence of the 1σ uncertainties in photo- z as a function of magnitude for $0.2 < z < 1.25$. The z_p for the CFHTLS-DEEP were derived from 5 broad bands (u^* , g' , r' , i' , z'). The accuracy of the COSMOS photo- z is improved by a factor of 3 when compared to CFHTLS-DEEP. This improvement is largely due to the 12 medium bands in this redshift range (NIR data have a real impact only at $z > 1.25$). We checked this by deriving photo- z without the 12 medium bands, obtaining an accuracy of $\sigma_{\Delta z/(1+z_s)} = 0.03$ at $i^+ < 22.5$ (similar to the COSMOS release of Mobasher et al. 2007 using 8 broad bands u^* , B_J , V_J , g^+ , r^+ , i^+ , z^+ and K_S). COMBO-17 includes 12 medium bands in addition to the 5 broad bands and Wolf et al.

(2004) achieved an accuracy of $\sigma_{\Delta z/(1+z)} = 0.02$ at $i^+ \sim 21.5$. The accuracies of COMBO-17 photo- z are intermediate between those of COSMOS-30 and CFHTLS for $i_{AB}^+ < 22.5$ but become larger that the CFHTLS accuracies at fainter magnitudes, because the COMBO-17 data are about 1.5 mag shallower than the CFHTLS data. Since only secure photo- z (one solution) have been kept in the COMBO-17 catalogue, it explains the flattening of the error bars at $i_{AB}^+ > 24$.

5.2. Redshift Distribution of Galaxies

Fig. 13 shows the galaxy redshift distribution per deg^2 from the COSMOS and CFHTLS surveys (for $21.5 < i_{auto}^+ < 24.5$). Although the surveys are largely independent and the photo- z are computed with different codes, the overall agreement in the redshift distributions is excellent. The agreement is particularly good between COSMOS and CFHTLS-D2 which covers 1 deg^2 within the COSMOS field.

The density of galaxies at $z > 1.5$ in the CFHTLS fields is of a crucial interest for weak

mag range	a	b	c	A	average z	median z
$22.0 < i_{auto}^+ < 22.5$	0.497 ± 0.019	12.643 ± 0.409	0.381 ± 0.016	4068.19	0.66	0.66
$22.5 < i_{auto}^+ < 23.0$	0.448 ± 0.016	9.251 ± 0.218	0.742 ± 0.030	9151.98	0.76	0.72
$23.0 < i_{auto}^+ < 23.5$	0.372 ± 0.012	6.736 ± 0.094	1.392 ± 0.055	18232.24	0.90	0.82
$23.5 < i_{auto}^+ < 24.0$	0.273 ± 0.008	5.281 ± 0.039	2.614 ± 0.096	35508.58	1.05	0.92
$24.0 < i_{auto}^+ < 24.5$	0.201 ± 0.005	4.494 ± 0.024	3.932 ± 0.134	60306.30	1.18	1.00
$24.5 < i_{auto}^+ < 25.0$	0.126 ± 0.003	4.146 ± 0.021	5.925 ± 0.191	103340.04	1.25	1.06

Table 2: Galaxy redshift distribution per deg^2 . The parameters a , b , c and A of the eqn.4 function (Fu et al. 2008) are given per apparent bin.




Fig. 11.— Same as Fig.8 for the CFHTLS-Deep survey (Ilbert et al. 2006) and the COMBO-17 survey (Wolf et al. 2004).

lensing analysis (Benjamin et al. 2007). The CFHTLS redshift distribution presents a bump at $z \sim 3$, but this excess could be due to misidentifications between the $z < 0.4$ and $z > 1.5$ photo- z (Van Waerbeke et al. 2008) when no NIR data are available (as is the case for CFHTLS). The COSMOS photo- z are computed with NIR data and such catastrophic failures are limited (see §4.1). Fig. 13 shows that this bump seems created by a deficit of galaxies at $2.5 < z < 3$, by comparing CFHTLS-D2 and COSMOS.




Fig. 12.— 1σ error for the z_p estimate as a function of the apparent magnitude in the redshift range $0.2 < z_p \leq 1.25$. The 1σ errors have been rescaled by a factor 1.2 for the COMBO-17 survey (Wolf et al. 2004) (see text and Fig. 11).

Fu et al. (2008) used the CFHTLS photo- z distribution to estimate the matter density parameter Ω_m and the amplitude of the matter power spectrum σ_8 . The redshift distribution from Fu et al. (2008) at $21.5 < i^+ < 24.5$ and $z < 2.5$ is overplotted in Fig. 13 with and without (J. Coupon, private communication) the weight applied for the weak lensing selection (dashed and dashed-dot lines, respectively). Given the 20% variation ex-




Fig. 13.— Redshift distribution from the 2-deg² COSMOS survey (black solid line), from the 1-deg² CFHTLS-D2 field (green dashed line), from the 4-deg² CFHTLS-DEEP fields (blue dotted line, including also D2). CFHTLS-D2 covers 1 deg² within the COSMOS field. The red long dashed line is the redshift distribution obtained by Fu et al. (2008) who fit the CFHTLS-DEEP photo-z in the magnitude bin $21.5 < i^+ < 24.5$ and the dashed-dotted line is obtained without the weight applied for the CFHTLS weak lensing selection (J. Coupon, private communication).

pected from cosmic variance (Ilbert et al. 2006), it is in excellent agreement with the COSMOS-30 redshift distribution. This agreement suggests that the derivation of $\sigma_8(\Omega_m/0.25)^{0.64} = 0.785 \pm 0.043$ by Fu et al. (2008) is not suffering from biases due to the photo-z or significant cosmic variance.

The COSMOS-30 redshift distribution can be fit with a parametrization similar to that used by Fu et al. (2008):

$$n(z) = A \frac{(z^a + z^{ab})}{(z^b + c)} \quad (4)$$

where A is the normalization factor and a , b and c are free parameters. The best fit parameters a , b and c are given in Table 2, as well as the




Fig. 14.— Top panel: Evolution of the redshift distribution as a function of i^+ magnitude in the COSMOS field. We use the parametrization of Fu et al. (2008) in different redshift bins. Bottom panel: Cumulative redshift distribution.

median redshifts. Fig. 14 shows the best fit redshift distribution per apparent magnitude bin. As expected, the median redshift increases at fainter apparent magnitude, ranging from $z_m = 0.66$ at $22 < i^+ < 22.5$ to $z_m = 1.06$ at $24.5 < i^+ < 25$.

6. Summary

This paper presents a new version of the photometric redshift catalog for the 2-deg² COSMOS survey computed with new ground-based NIR data, deeper IRAC data and a new set of 12 medium bands from the Subaru Telescope. The COSMOS photometry now includes a total of 30 filters – from the UV (*GALEX*) to the MIR (*Spitzer*-IRAC). The photo-z catalogue derived here contains 607,617 sources at $i^+ < 26$. The 1887 *XMM*-COSMOS sources (mainly AGN) are not included in this catalogue; their photo-z are derived in Salvato et al. (2008) with similarly good accuracy using a set of templates for composite AGN/galaxies.

The galaxy photo-z were tested and improved using spectroscopic redshift samples from the

Sample		N	median z_s	median i^+	$\sigma_{\Delta z/(1+z_s)}$	η in %
zCOSMOS bright	$17.5 < i^+ < 22.5$	4146	0.48	21.6	0.007	0.7
zCOSMOS faint	$1.5 < z_s < 3$	147	2.20	24.0	0.054	20.4
MIPS bright	$17.5 < i^+ < 22.5$	186	0.68	21.7	0.009	0.0
MIPS faint	$22.5 < i^+ < 24.0$	116	0.90	23.1	0.011	0.0
MIPS very faint	$24.0 < i^+ < 25.0$	15	1.15	24.2	0.053	20.0

Table 3: Redshift accuracies estimated from the comparison between photometric redshifts and spectroscopic redshifts.

zCOSMOS and Keck surveys. Biases in the photo-z were removed by iterative calibration of the photometric band zero-points. As suggested by the data, two different dust extinction laws were applied specific to the different SED templates. A new method to take into account the emission lines was implemented using relations between the UV continuum and the emission line fluxes associated with star formation activity. The allowance for emission lines decreased the photo-z dispersion by a factor 2.5.

Based on a comparison between our new values for z_p and 4148 measured values of z_s from zCOSMOS, we estimate an accuracy of $\sigma_{\Delta z/(1+z_s)} = 0.007$ for the galaxies brighter than $i^+ = 22.5$. The accuracies measured with the various spectroscopic samples are summarized in table.3. We extrapolate this result to fainter magnitudes using the 1σ uncertainties in the photometric redshift probability functions. This is found to provide reliable uncertainty estimates for the photo-z technique developed here since these uncertainties agree with the dispersion in the offsets of photo-z from spec-z for the spectroscopic sample. At $z < 1.25$, we estimate a photo-z accuracy of $\sigma_{\Delta z} = 0.02, 0.04, 0.07$ for $i^+ \sim 24, i^+ \sim 25, i^+ \sim 25.5$. The accuracy is strongly degraded at $i^+ > 25.5$ and the exploitation of the COSMOS-30 photo-z at fainter magnitudes should be done carefully. The accuracy is 3-5 times better than the photo-z determined for the CFHTLS-DEEP (Ilbert et al. 2006) and the previous COSMOS photo-z release (Mobasher et al. 2007), and 2 times better than the accuracy of the photo-z determined for COMBO-17 at $i^+ < 23$ (Wolf et al. 2004). Deep NIR (J, K) and IRAC data were essential to keep the catastrophic failures low at

$z > 1.25$. We note that the accuracy of the COSMOS photo-z could soon be further improved at $z > 1.25$ with the addition of new data currently being obtained in the Y band (UKIRT), H band (CFHT) and the ULTRA-VISTA survey.

Our photo-z catalogue contains 607,617 sources at $i^+ < 26$. The accurate photo-z derived here for this extremely large sample of galaxies are crucial to scientifically exploit the full legacy value of the multi- λ data sets in the COSMOS field (*HST*-ACS, *Spitzer*, *GALEX*, VLA, *XMM* and *Chandra*).

We gratefully acknowledge the contributions of the entire COSMOS collaboration consisting of more than 100 scientists. The *HST* COSMOS program was supported through NASA grant HST-GO-09822. French co-authors acknowledge support from the French Agence National de la Recherche fund ANR-07-BLAN-0228 (DESIR project). More information on the COSMOS survey is available at <http://www.astro.caltech.edu/cosmos>. We also greatly appreciate the hospitality provided by the Aspen Center for Physics where this manuscript was completed.

REFERENCES

- Arnouts, S. et al. 1999, MNRAS, 310, 540
- Arnouts, S., Walcher, C. J., Le Fèvre O. et al. 2007, A&A, 476, 137
- Ball N.M., Loveday J., Fukugita M., Nakamura O., Okamura S., Brinkmann J. & Brunner R.J., 2004, MNRAS, 348, 1038
- Baum, W.A. 1962, Problems of Extra-Galactic Research, 15, 390

- Benjamin J., Heymans C., Semboloni E. et al. 2007, MNRAS, 381, 702
- Bertin E. & Arnouts S. 1996, A&AS, 117, 393
- Bixler J.V., Bowyer S. & Laget M., A&A, 250, 370
- Bolzonella, M., Miralles, J.-M. & Pelló, R. 2000, A&A, 363, 476
- Boulade, O., Charlot, X., Abbon, P. et al. 2003, SPIE, 4841, 72
- Brodwin, M., Lilly, S.J., Porciani, C. et al. 2006, ApJS, 162, 20
- Brusa, M., Zamorani, G., Comastri, A. et al. 2007, ApJS, 172, 353
- Bruzual, G., & Charlot, S. 2003, MNRAS, 344, 1000
- Budavári, T., Szalay, A. S., Connolly, A. J., Csabai, I. & Dickinson, M. 2000, AJ, 120, 1588
- Calzetti, D., Armus, L., Bohlin, R., C. et al. 2000, ApJ, 533, 682
- Capak, P. et al., 2008, in preparation
- Capak, P., Aussel, H., Ajiki, M. et al. 2007, ApJS, 172, 99
- Caputi K.I., McLure R.J., Dunlop J.S., Cirasuolo M. & Schael A.M., 2006, MNRAS, 366, 609
- Caputi K.I., Lilly S.J., Aussel H. et al., 2008, ApJ, 680, 939
- Chabrier G., Baraffe I., Allard F. & Hauschildt P., 2000, ApJ, 542, 464
- Coleman, G. D., Wu, C.-C., Weedman, D.W. 1980, ApJS, 43, 393
- Collister, A. A. & Lahav, O. 2004, PASP, 116, 345
- Daddi, E., Cimatti, A., Renzini, A. et al. 2004, ApJL, 600, 127
- Feldmann, R., Carollo, C. M., Porciani, C. et al. 2006, MNRAS, 372, 565
- Fernández-Soto, A., Lanzetta, K. M. & Yahil, A. 1999, ApJ, 513, 34
- Fischera, J., Dopita, M. A., & Sutherland, R. S. 2003, ApJL, 599, L21
- Fontana, A., D’Odorico, S., Poli, F. et al. 2000, AJ, 120, 2206
- Fu, L., Semboloni, E., Hoekstra, H. et al. 2008, A&A, 479, 9
- Gabasch, A., Bender, R., Seitzer, S. al. 2004, A&A, 421, 41
- Guillaume M., Llebaria A., Aymeric D., Arnouts S. & Milliard B. 2006, SPIE, 6064, 332
- Heinis, S., Milliard, B., Arnouts, S. et al. 2007, ApJS, 173, 503
- Hoaglin, D. C., Mosteller, F., & Tukey, J. W 1983, Understanding Robust and Exploratory Data Analysis. John Wiley & Sons, 404-414.
- Ilbert, O., Arnouts, S., McCracken, H. J. et al. 2006, A&A, 457, 841
- Kartaltepe, J. et al. 2008, in preparation
- Kennicutt, R.C. 1998, ARA&A, 36, 189
- Lamareille, F. et al. 2008, in preparation
- Leauthaud, A., Massey, R., Kneib, J.-P. et al. 2007, ApJS, 172, 219
- Le Fèvre, O., Vettolani, G., Paltani, S. et al. 2004, A&A, 428, 1043
- Le Fèvre, O., Vettolani, G., Garilli, B. et al. 2005, A&A, 439, 845
- Lilly S. J., Le Fèvre, O., Renzini, A. et al. 2007, ApJS, 172, 70
- Lilly, S. J. et al., 2008, in preparation
- Lonsdale, C. J., Smith, H. E., Rowan-Robinson, M. et al. 2003, PASP, 115, 897
- Madau, P., Ferguson, H. C., Dickinson, M. E. et al. 1996, MNRAS, 283, 1388
- Maraston, C., Daddi, E., Renzini, A., Cimatti, A. et al. 2006, ApJ, 652, 85
- Mazure, A., Adami, C., Pierre, M. et al. 2007, A&A, 467, 49
- McCall, M. L., Rybski, P. M., & Shields, G. A. 1985, ApJS, 57, 1

- McCracken, H. J., Ilbert, O., Mellier, Y. 2007, A&A, 479, 431
- McCracken, H. J. et al. 2008, in preparation
- Mobasher, B., Capak, P., Scoville, N. Z. et al. 2007, ApJS, 172, 117
- Mouhcine, M., Lewis, I., Jones, B., Lamareille, F., Maddox, S. J. & Contini, T. 2005, MNRAS, 362, 1143
- Moustakas, J., Kennicutt, R. C. & Tremonti, C. A. 2006, ApJ, 642, 775
- Noll, S., Pierini, D., Pannella, M. & Savaglio, S. 2007, A&A, 472, 455
- Peacock, J. A., Schneider, P., Efstathiou, G. et al. 2006, ESA-ESO Working Group on "Fundamental Cosmology", arXiv:astro-ph/0610906
- Polletta, M., Tajer, M., Maraschi, L. et al. 2007, ApJ, 663, 81
- Prevot, M. L., Lequeux, J., Prevot, L., Maurice, E. & Rocca-Volmerange, B. 1984, A&A, 132, 389
- Rowan-Robinson, M., Babbedge, T., Oliver, S. et al. 2008, MNRAS, 386, 697
- Sanders, D. B., Salvato, M., Aussel, H. et al. 2007, ApJS, 172, 86
- Sasaki, S. et al. 2008, ApJ, in preparation
- Salvato, M. et al. 2008, ApJ, in preparation
- Sawicki, M. J., Lin, H., Yee, H. K. C. 1997, AJ, 113, 1
- Scoville, N. Z., Abraham, R. G., Aussel, H. et al. 2007, ApJS, 172, 38
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525
- Silva, L., Granato, G. L., Bressan, A. et al. 1998, ApJ, 509, 103
- Steidel, C. C., Shapley, A., Pettini, M. et al. 2004, ApJ, 604, 534
- Surace, J. A., Shupe, D. L., Fang, F. 2004, VizieR Online Data Catalog, 2255, 0
- Taniguchi, Y., Scoville N. Z., Murayama, T. et al. 2007, ApJS, 172, 9
- Taniguchi, Y. et al. 2008, ApJ, in preparation
- van Waerbeke, L. 2008, ApJ, in preparation
- Wang, L., & Steinhardt, P. J. 1998, ApJ, 508, 483
- Wolf C., Meisenheimer, K., Rix, H.-W. et al. 2003, A&A, 401, 73
- Wolf C., Meisenheimer, K., Kleinheinrich, M. et al. 2004, A&A, 421, 913
- Zamojski, M. A., Schiminovich, D., Rich, R. M. et al. 2007, ApJS, 172, 468
- Zucca, E., Ilbert, O., Bardelli, S. et al. 2006, A&A, 455, 879